Journal of Zhejiang University-SCIENCE B (Biomedicine & Biotechnology) ISSN 1673-1581 (Print); ISSN 1862-1783 (Online) www.zju.edu.cn/jzus; www.springerlink.com E-mail: jzus@zju.edu.cn



Fungal spondylodiscitis in a patient recovered from H7N9 virus infection: a case study and a literature review of the differences between *Candida* and *Aspergillus* spondylodiscitis*#

Lie-dao YU[§], Zhi-yun FENG[§], Xuan-wei WANG, Zhi-heng LING, Xiang-jin LIN^{†‡}

(Department of Orthopedic Surgery, the First Affiliated Hospital, School of Medicine, Zhejiang University, Hangzhou 310003, China) [†]E-mail: doclinxiangjin@163.com

Received Feb. 24, 2016; Revision accepted June 23, 2016; Crosschecked Oct. 18, 2016

Abstract: To report a rare case of fungal spondylodiscitis in a patient recovered from H7N9 virus infection and perform a literature review of the different characteristics of Candida and Aspergillus spondylodiscitis, we reviewed cases of spondylodiscitis caused by Candida and Aspergillus species. Data, including patients' information, pathogenic species, treatment strategy, outcomes, and relapses, were collected and summarized. The characteristics of Candida and Aspergillus spondylodiscitis were compared to see if any differences in clinical features, management, or consequences could be detected. The subject of the case study was first misdiagnosed as having a vertebral tumor, and then, following open biopsy, was diagnosed as having fungal spondylodiscitis. The patient made a good recovery following radical debridement. Seventy-seven additional cases of Candida spondylodiscitis and 94 cases of Aspergillus spondylodiscitis were identified in the literature. Patients with Candida spondylodiscitis tended to have a better outcome than patients with Aspergillus spondylodiscitis (cure rate 92.3% vs. 70.2%). Candida was found more frequently (47.8%) than Aspergillus (26.7%) in blood cultures, while neurological deficits were observed more often in patients with Aspergillus spondylodiscitis (43.6% vs. 25.6%). Candida spinal infections were more often treated by radical debridement (60.5% vs. 39.6%). Patients with Candida spondylodiscitis have better outcomes, which may be associated with prompt recognition, radical surgical debridement, and azoles therapy. A good outcome can be expected in fungal spondylodiscitis with appropriate operations and anti-fungal drugs.

Key words: Spondylodiscitis, Spinal infection, H7N9 virus, Avian influenza virus, *Candida*, *Aspergillus*, Fungi, Review http://dx.doi.org/10.1631/jzus.B1600077 **CLC number:** R681.5+1

1 Introduction

Invasive fungal spinal infection is a rare condition, compared with pyogenic or tuberculous spondylodiscitis (Broner *et al.*, 1996; Chia *et al.*, 2005; D'Agostino *et al.*, 2010). It often occurs as an op-

portunistic infection in immunocompromised patients, but immunocompetents can also be affected. Back pain is the most frequent manifestation, but neurological deficits may sometimes occur (Kim *et al.*, 2006). Diagnosis of fungal spondylodiscitis without a biopsy or operation is very difficult, due to the non-specific symptoms, the low rate of positive blood cultures, and atypical radiological findings.

The optimal management of fungal spondy-lodiscitis remains uncertain. Several reports suggest that fungal spondylodiscitis without neurological deficits and spinal instability can be treated merely by anti-fungal drugs (Sugar *et al.*, 1990; Cortet *et al.*, 1994; Hennequin *et al.*, 1996; Rachapalli *et al.*, 2010;

[‡] Corresponding author

[§] The two authors contributed equally to this work

^{*} Project supported by the Zhejiang Medical Technology and Education Foundation (No. 2015107339), China

[#] Electronic supplementary materials: The online version of this article (http://dx.doi.org/10.1631/jzus.B1600077) contains supplementary materials, which are available to authorized users

ORCID: Xiang-jin LIN, http://orcid.org/0000-0003-0062-0448

Zhejiang University and Springer-Verlag Berlin Heidelberg 2016

Ersoy et al., 2011; Theodoros and Sotirios, 2012). Others have argued that surgical treatments should be given as soon as there is no clear diagnosis and when conventional treatment has failed, and should not be limited to cases with neurological presentations and spinal instabilities (Nasca and McElvein, 1985; Bridwell et al., 1990; Frazier et al., 2001; Hendrickx et al., 2001; Lenzi et al., 2004; Skaf et al., 2010; Iwata et al., 2014). Since a delayed operation may lead to a poor outcome (Iwata et al., 2014), many doctors recommend prompt surgical intervention. However, answers to some important questions about surgery for fungal spondylodiscitis are still unclear, such as how debridement should be given and how to choose drugs postoperatively.

As Aspergillus and Candida are reported to be two main etiologies for fungal spinal infections (Skaf et al., 2010), a literature review of spinal infections caused by Aspergillus and Candida may be helpful to clarify these questions. Also, there are no reports regarding the differences between the characteristics of Candida and Aspergillus spondylodiscitis. The virulence of Aspergillus has been shown to be stronger than that of Candida. Furthermore, Aspergillus infections in other systems have high mortality rates. It is unclear whether the consequences of Candida and Aspergillus spondylodiscitis are different, or if there should be any differences in their management. These are important issues, as they affect the decision making of doctors when confronted with fungal spondylodiscitis.

Here, we report a case of *Candida* spondylodiscitis in a patient recovered from H7N9 infection. We also review the characteristics of spondylodiscitis caused by *Candida* and *Aspergillus* described in previous reports, in the hope that this can provide insights for the management of fungal spinal infections.

2 Case presentation

In April 2013, a 76-year-old man, with a past medical history of H7N9 virus infection, presented with progressive lumbar pain for three months. On April 11, 2013, he had a heavy cough and was diagnosed at our hospital as having an H7N9 infection (Gao *et al.*, 2013). The patient had severe pulmonary inflammation due to the H7N9 infection (Fig. 1). He

was transferred to the intensive care unit, and antivirus and glucocorticoid treatments were given immediately (Gao et al., 2013). After one month, the patient showed a good clinical progression and was discharged. Three months later, he complained of pain in the lower back and left leg, accompanied by a low fever. The pain was unbearable and was aggravated by a change of position. Magnetic resonance imaging (MRI) showed signal changes at the L5 and S1 vertebral body, without disc involvement or any surrounding abscess (Figs. 2a-2c). Emission computed tomography indicated a concentration of radioactivity on the L5 vertebra. Therefore, a spinal tumor was first considered and a biopsy through the vertebral pedicle was performed. Pathological tests showed that the sample consisted of inflammatory fibroid tissue with active T lymphocyte proliferation, but there was no sign of tumor.




Fig. 1 Computed tomographic scan of the chest of the patient

After the onset of infection by H7N9, consolidations were observed in almost the whole right lung and ground-glass opacities were found in the left lung

During this period, little pain relief was achieved by pain killers, and the low fever continued. Another MRI scan revealed signal changes in the L5 and S1 endplate with the disc involved, indicating spondylodiscitis at the L5/S1 level (Figs. 2d–2f). At this time, lumbar spine tuberculosis (TB) was considered. The patient underwent conventional anti-TB therapy with isoniazid, rifampicin, ethambutol, and moxifoxacin. However, two months of chemotherapy for TB produced no improvement. Based on the limited response to anti-TB therapy, we realized that TB might not be the pathogen.




Fig. 2 Lumbar magnetic resonance images (MRIs) of the nation

MRI at the first time showed hypo-intensity on the whole L5 vertebrae body and the posterior endplate of S1 on the sagittal T1 weighted (T1W) image (a), while hyper-intensity can be observed only at the endplate of L5 on the sagittal T2 weighted (T2W) image (b). Mild signal changes can also be observed on the lateral image (c). Repeated MRI two months later showed that the area of hypo-intensity had widened, with the S1 vertebra body and the adjacent disc involved on the sagittal T1W image (d). Also, there were increased signal changes at the L5/S1 disc and adjacent endplates on the sagittal T2W image with both endplates having "moth-eaten" damage (e). The lateral T2W image showed erosion at the center of S1 vertebra (f)

The patient was then transferred to our department, as the pain was aggravated and accompanied by ache and numbness in the left lower limb. An exploration of the L5/S1 disc through the posterior medial approach was recommended to determine the source of the problem and also to perform a debridement. At surgery, the L5/S1 intervertebral disc and adjacent vertebral bodies appeared damaged. There was no abscess or pus, but a mass of granulation tissue was visible. The infected tissue with adjacent endplates was resected radically. An auto-bone graft and interbody fixation were inserted in the disc space. A direct smear and culture of the disc lesions showed no bacteria or mycotic organisms, but grew Candida albicans. Histologically, there was chronic inflammatory granuloma formation, necrosis, and fungal spores. Therefore, the patient received fluconazole intravenously for 1 month and then orally for 3 months. One year postoperatively, he was without back pain.

3 Materials and methods

PubMed and Google Scholar were searched using the terms spondylodiscitis, spondylitis, vertebral,

diskitis, disc, fungal, fungi, Aspergillus, and Candida. The language was restricted to English, and the species to humans. No other limit was used. Cases of spondylodiscitis caused by Candida and Aspergillus species were selected. References cited by these reports were thoroughly screened to find additional cases. All the data, including basic information (age, gender, complaints, fever or not, blood culture findings, neurological deficits, infection sites), pathogenic species, treatment strategy, outcomes, and relapses were recorded if stated unambiguously.

Specimens had to be confirmed by pathological tests or cultures from infected tissues. Anti-fungal drugs before and after surgery were recorded separately. The treatment strategy was classified as antifungal only, anti-fungal followed by surgery, or surgery followed by anti-fungal. The surgical options were classified as simple debridement (including laminectomy and debridement without fusion) and radical debridement (radical debridement with fusion). The drug selections postoperatively were classified as amphotericin B (amB) only, azoles only, amB+fluorocytosine, amB+azoles, amB+fluorocytosine+azoles, or micafungin/caspofungin.

4 Results

One hundred and seventy-one cases of *Candida* and *Aspergillus* spinal infections were identified in the literature (supplementary material). The characteristics of patients, including our patient, are summarized in Table 1. The mean age was 52.8 years (range, 11–89 years). Most of the patients were men (70.9%).

4.1 Pathogenesis

Among the 172 cases, 78 patients (45.3%) had documented *Candida* spinal infection, while 94 (54.7%) had *Aspergillus* spondylodiscitis (Table 1). The fungi were not commonly found in blood samples: only 55 (32.0%) cases showed positive findings, 41.0% of *Candida* spondylodiscitis and 24.5% of *Aspergillus* spondylodiscitis cases (Table 1).

4.2 Clinical presentation

Back pain was the main complaint associated with fungal spinal infection. Almost all the patients suffered from back pain to variable degrees. Fever was not commonly seen (31.4%), with 35.9% in

Candida spondylodiscitis and 27.7% in Aspergillus spondylodiscitis cases (Table 1). Neurological deficits were observed in 61 patients (35.5%), and were present more often in Aspergillus spondylodiscitis (43.6% vs. 25.6%) (Table 1). In 109 patients (63.4%), lumbar vertebrae were solely involved, and in 42 patients (24.4%), the thoracic spine was solely involved. The cervical spine was solely involved in only 2 patients with Candida spondylodiscitis. Multiple spinal infections were often caused by Aspergillus, with lumbar and thoracic spinal regions being the main concomitant sites (14 cases, 8.1%) (Table 1).

Table 1 Characteristics of patients with *Aspergillus* and *Candida* spinal infections

•		
Characteristics	Candida (n=78)	Aspergillus (n=94)
Age	55.2±18.0	46.4±15.1
Gender		
Male	58 (74.4%)	64 (68.1%)
Location		
C	2 (2.6%)	0
T	20 (25.6%)	22 (23.4%)
L	51 (65.4%)	58 (61.7%)
C+T	0	3 (3.2%)
T+L	5 (6.4%)	9 (9.6%)
C+T+L	0	2 (2.1%)
Fever	28 (35.9%)	26 (27.7%)
Pain	78 (100%)	94 (100%)
Neurological deficit	20 (25.6%)	41 (43.6%)
Blood culturea	32 (41.0%)	23 (24.5%)
Operation	44 (56.4%)	65 (69.1%)
Outcome ^b	72 (92.3%)	66 (70.2%)
Relapse	7 (9.0%)	8 (8.5%)

C: cervical spine; T: thoracic spine; L: lumbar spine. ^a 31 cases did not refer to data of blood culture (17 for *Candida* and 14 for *Aspergillus*); ^b 6 cases did not have data of outcome (each three for *Candida* and *Aspergillus*). Data were present as number of cases (percentage)

4.3 Treatments

Sixty-two of the 172 patients (36.0%) received only drug therapy, 54 (31.4%) required an operation after the failure of drug therapy, and 56 (32.6%) had an early operation combined with anti-fungal drugs (Table 2).

As for the surgical plan, 47 patients received simple debridement or laminectomy. The other 44 patients were treated with radical excision and bone grafts. For 19 patients, no surgical information was available (Table 3).

Table 2 Number of patients with *Candida* and *Aspergillus* spondylodiscitis receiving conservative or operative treatment

Treatment strategy	Candida	Aspergillus
Conservative treatment		
Anti-fungals only	34	28
Operative treatment		
Anti-fungals failed,	20	35
then operation		
Operation directly, then	24	31
anti-fungals		

Table 3 Surgical options for *Candida* and *Aspergillus* spondylodiscitis

Surgical strategy	Candida	Aspergillus
Simply debridement ^a	15	32
Radical debridement ^a	23	21
Approach (A:P:A+P) ^b	13:17:0	14:34:3
Stage (1:2)	30:1	48:4

^a Twelve cases did not refer to the detailed information about the operation way (6 for *Candida*, 13 for *Aspergillus*). ^b Twentynine cases did not refer to the approach of operations (14 for *Candida* and 15 for *Aspergillus*). In addition, transthoracic and extrapleural anterolateral approaches were classified as anterior approach, and costotransversectomy was classified as posterior approach for surgery in the thoracic spine. A, anterior; P, posterior; A+P, anterior plus posterior

Post-operative treatment information was available for 97 of the 110 patients who underwent surgery. Azoles were more likely to be used by patients with *Candida* spinal infection (19/40 vs. 19/57), and amB by patients with *Aspergillus* spinal infection (25/57 vs. 7/40) (Table 4). Combinations of amB/azoles (16/97), amB/fluorocytosine (5/97), and amB/fluorocytosine/azoles (3/97) were not usually used after the surgery. New drugs, such as micafungin and caspofungin, were not commonly used after surgery: only three patients infected by *Candida* were given micafungin.

Table 4 Drug selections after operations

O	•	
Drug	Candida	Aspergillus
AmB only	7	25
AmB+fluorocytosine	2	3
Azoles only	19	19
AmB+azoles	7	9
AmB+fluorocytosine+azole	2	1
Micafungin/caspofungin	3	0

Azole includes voriconazole, fluconazole, itraconazole, miconazole, ketoconazole, cotrimoxazole, and posaconazole. AmB indicates amphotericin B. Four cases in *Candida* spinal infection and 9 cases in *Aspergillus* spinal infection have no data about the drug therapy post-operatively

4.4 Outcomes

Outcomes could not be evaluated for six patients because of insufficient information. With the exception of two patients, all patients with *Candida* spinal infections were ultimately cured by means of medicine and/or surgical treatment. The cure rate for *Candida* spondylodiscitis (92.3%) was higher than that for *Aspergillus* spondylodiscitis (70.2%) (Table 1). Fifteen patients (8.7%) suffered relapses. The relapse rate of *Candida* spondylodiscitis (9.0%) paralleled that of *Aspergillus* spondylodiscitis (8.5%) (Table 1).

5 Discussion

The pathogenic organisms of spondylodiscitis are often bacteria or Mycobacterium tuberculosis, while fungi are seldom implicated. However, the incidence of fungal spinal infections has increased in recent years, associated with the widespread use of broad-spectrum antibiotics, prolonged chemotherapeutic drugs, and corticosteroids (Kim et al., 2006; Shi et al., 2015). To our knowledge, this is the first recorded case of fungal spondylodiscitis found in a patient recovered from H7N9 infection. As outbreaks of H7N9 and other avian influenza virus infections increase worldwide (Liem et al., 2009; Yang et al., 2012; Chen et al., 2013; Gao et al., 2013; Uyeki and Cox, 2013), we should keep in mind that patients infected by such viruses may suffer from fungal spondylodiscitis, even after recovery from the viral infection.

Heavy corticosteroid consumption is the main risk factor for these opportunistic infections. Also, the viral infection may facilitate the proliferation of other pathogens growing in the blood, which may then seed in the vertebral body (Ceroni et al., 2013; 2014). Previous studies have shown that early viral infection can damage the mucosal layer of the airway, leading to respiratory tract bacterial infections (Yagupsky, 2004). A limitation of our study was that we did not check for H7N9 viral infection in the disc by polymerase chain reaction (PCR) or other assays to determine whether H7N9 might be involved in the pathogenesis of Candida spondylodiscitis. A deficiency of the immune system may be another factor predisposing for the invasion of Candida in the vertebral body. It is well accepted that viral infection,

corticosteroid consumption, and prolonged periods of intensive care will cause temporary depression of the immune function, making patients susceptible to various types of infections.

Aspergillus and Candida have been shown to be the predominant causative fungi of spinal infections (Kim et al., 2006; Skaf et al., 2010), consistent with the findings of our review. Other species of fungi, such as Blastomyces, Pseudallescheria boydii, and Scremonium, are only rarely pathogens of spinal infections (Groll and Walsh, 2001; Kim et al., 2006; Skaf et al., 2010). In our study, the cure rate of Candida spinal infection was higher than that of Aspergillus (92.3% vs. 70.2%). This may be attributable to the difference in virulence between these pathogens. Systematic Aspergillus infection is life-threatening and has a high mortality rate (Mawk et al., 1983; Walsh, 2008; Pappas et al., 2009). Multiple vertebral infections by Aspergillus, as demonstrated by our review and Kwon et al. (2011), may also contribute to the lower cure rate of Aspergillus spinal infections. Early recognition of *Candida* spinal infection may also affect the different outcomes, since we found that patients with Candida spondylodiscitis had a higher positive rate of blood cultures. A more sensitive test is needed for detecting Aspergillus in blood cultures, because blood cultures are always the first step in testing patients with suspected infections.

In this review, only 36.0% of patients were treated by antifungal drugs alone. Such treatment often fails to stop the progression of bone destruction and can necessitate subsequent surgical intervention. Surgery was required following antifungal treatment in 32.0% of cases in this review and 30.0% of cases in another study (Hendrickx et al., 2001). The majority of patients were treated with surgical debridement, either to remove infected tissue, or to enable a definitive diagnosis, while indications for surgery varied from reports. Generally, open surgery could be divided into simple debridement and radical debridement with internal fixation. In this review, patients with Candida spondylodiscitis often underwent radical debridement, while simple debridement was chosen more frequently by patients with Aspergillus spondylodiscitis. Radical debridement can relieve the burden of infection and enhance the infiltration of antifungal drugs into the infected vertebra and disc. As Aspergillus spinal infection has a higher rate of mortality, a radical excision of the infected tissue is recommended if the condition of the patient permits. If not, patients with minimal bone destruction and a small abscess can be treated by endoscopic surgery in the early phase, without causing progressive instability (Iwata *et al.*, 2014).

Even after surgical removal of infected tissue, sufficient anti-fungal treatment is very important. We collected information about drug selection post-operatively in the case reports to determine implications for clinical practice. AmB and azoles, used solely or in combination, were the basic regimens for the treatment of fungal spondylodiscitis. New drugs, such as caspofungin and micafungin, were always used when azole-resistance or amB-resistance occurred (Pemán *et al.*, 2006; Theodoros and Sotirios, 2012; Tan *et al.*, 2014). As the course of treatment varied from 1 to 24 months, these data were not analyzed. In general, amB was given for 2–8 weeks depending on the tolerance and response of patients, while azoles were always given for 3–24 months.

The key factor differentiating spondylodiscitis from a vertebral tumor in MRIs is the involvement of the disc, as the pathogens invade the endplates leading to infection of the disc. The initial MRIs in our case, however, showed no signal changes in the adjacent disc, consistent with the MR findings of fungal spondylodiscitis in previous studies (Williams *et al.*, 1999; Shashidhar *et al.*, 2014; Storm *et al.*, 2014). This explains why our case was at first misdiagnosed as a vertebral tumor. Fungal spinal infections in the early phase are insidious, and MRI scans can mimic the presentation of a vertebral tumor.

The insidious clinical presentations, low frequency of positive blood cultures, non-specific radiological findings and poor awareness add to the difficulty in diagnosing fungal spinal infection (Frazier *et al.*, 2001; Kim *et al.*, 2006; Skaf *et al.*, 2010). Biopsy or open surgery should be performed immediately when anti-bacterial or anti-TB treatments fail to control the symptoms. However, biopsy sometimes cannot provide an accurate answer (Palmisano *et al.*, 2011; Shashidhar *et al.*, 2014). A non-diagnostic aspirate should prompt an open biopsy for pathogen cultures and pathological examinations. Culture samples should be evaluated for not only bacteria and mycobacterial organisms but also fungi, if spondylodiscitis are suspected.

In conclusion, we should keep in mind that fungal spondylodiscitis can be a complication in patients with avian influenza virus infection. *Candida* spondylodiscitis tends to have a better outcome than *Aspergillus* spondylodiscitis, maybe due to earlier detection, radical debridement, and lengthy azoles treatment. With regard to treatment, the present case had a good result following one-stage radical debridement and internal fixation.

Compliance with ethic guideline

Lie-dao YU, Zhi-yun FENG, Xuan-wei WANG, Zhi-heng LING, and Xiang-jin LIN declared that they have no conflict of interest.

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2008 (5). Informed consent was obtained from the patient for being included in the study and the work was approved by the ethical committee of the First Affiliated Hospital, School of Medicine, Zhejiang University (Hangzhou, China).

References

Bridwell, K., Campbell, J.W., Barenkamp, S.J., 1990. Surgical treatment of hematogenous vertebral *Aspergillus* osteomyelitis. *Spine*, **15**(4):281-285.

http://dx.doi.org/10.1097/00007632-199004000-00006

Broner, F.A., Garland, D.E., Zigler, J.E., 1996. Spinal infections in the immunocompromised host. *Orthop. Clin. N. Am.*, **27**(1):37-46.

Ceroni, D., Dubois-Ferrière, V., Cherkaoui, A., *et al.*, 2013. 30 years of study of *Kingella kingae*: post tenebras, lux. *Future Microbiol.*, **8**(2):233-245. http://dx.doi.org/10.2217/fmb.12.144

http://dx.doi.org/10.221//hiib.12.144

- Ceroni, D., Kampouroglou, G., Valaikaite, R., et al., 2014. Osteoarticular infections in young children: what has changed over the last years? Swiss Med. Wkly., 144: w13971.
- Chen, Y., Liang, W.F., Yang, S.G., *et al.*, 2013. Human infections with the emerging avian influenza A H7N9 virus from wet market poultry: clinical analysis and characterisation of viral genome. *Lancet*, **381**(9881):1916-1925. http://dx.doi.org/10.1016/S0140-6736(13)60903-4
- Chia, S.L., Tan, B.H., Tan, C.T., *et al.*, 2005. *Candida* spondylodiscitis and epidural abscess: management with shorter courses of anti-fungal therapy in combination with surgical debridement. *J. Infect.*, **51**(1):17-23. http://dx.doi.org/10.1016/j.jinf.2004.08.020
- Cortet, B., Richard, R., Deprez, X., et al., 1994. Aspergillus spondylodiscitis: successful conservative treatment in 9 cases. J. Rheumatol., 21(7):1287-1291.
- D'Agostino, C., Scorzolini, L., Massetti, A.P., et al., 2010. A seven-year prospective study on spondylodiscitis:

- epidemiological and microbiological features. *Infection*, **38**(2):102-107.
- http://dx.doi.org/10.1007/s15010-009-9340-8
- Ersoy, A., Dizdar, O.S., Koc, A.O., 2011. *Aspergillus fumigatus* spondylodiskitis in renal transplant patient: voriconazole experience. *Exp. Clin. Transplant.*, **9**(4): 265-269.
- Frazier, D.D., Campbell, D.R., Garvey, T.A., *et al.*, 2001. Fungal infections of the spine. Report of eleven patients with long-term follow-up. *J. Bone Joint Surg. Am.*, **83-A**(4):560-565.
- Gao, H.N., Lu, H.Z., Cao, B., *et al.*, 2013. Clinical findings in 111 cases of influenza A (H7N9) virus infection. *N. Engl. J. Med.*, **368**(24):2277-2285. http://dx.doi.org/10.1056/NEJMoa1305584
- Groll, A.H., Walsh, T.J., 2001. Uncommon opportunistic fungi: new nosocomial threats. *Clin. Microbiol. Infect.*, 7(Suppl. 2): 8-24.
 - http://dx.doi.org/10.1111/j.1469-0691.2001.tb00005.x
- Hendrickx, L., Wijngaerden, E.V., Samson, I., et al., 2001. Candidal vertebral osteomyelitis: report of 6 patients, and a review. Clin. Infect. Dis., 32(4):527-533. http://dx.doi.org/10.1086/318714
- Hennequin, C., Bourée, P., Hiesse, C., *et al.*, 1996. Spondylodiskitis due to *Candida albicans*: report of two patients who were successfully treated with fluconazole and review of the literature. *Clin. Infect. Dis.*, **23**(1):176-178. http://dx.doi.org/10.1093/clinids/23.1.176
- Iwata, A., Ito, M., Abumi, K., et al., 2014. Fungal spinal infection treated with percutaneous posterolateral endoscopic surgery. J. Neurol. Surg. Part A: Cent. Eur. Neurosurg., 75(3):170-176. http://dx.doi.org/10.1055/s-0032-1329268
- Kim, C.W., Perry, A., Currier, B., et al., 2006. Fungal infections of the spine. Clin. Orthop. Relat. Res., 444:92-99. http://dx.doi.org/10.1097/01.blo.0000203451.36522.4c
- Kwon, J.W., Hong, S.H., Choi, S.H., et al., 2011. MRI findings of Aspergillus spondylitis. Am. J. Roentgenol., 197(5): W919-W923. http://dx.doi.org/10.2214/AJR.11.6786
- Lenzi, J., Agrillo, A., Santoro, A., et al., 2004. Postoperative spondylodiscitis from *Aspergillus fumigatus* in immunocompetent subjects. *J. Neurosurg. Sci.*, **48**(2):81-85.
- Liem, N.T., Tung, C.V., Hien, N.D., 2009. Clinical features of human influenza A (H5N1) infection in Vietnam: 2004–2006. *Clin. Infect. Dis.*, **48**(12):1639-1646. http://dx.doi.org/10.1086/599031
- Mawk, J.R., Erickson, D.L., Chou, S.N., *et al.*, 1983. *Aspergillus* infections of the lumbar disc spaces. Report of three cases. *J. Neurosurg.*, **58**(2):270-274. http://dx.doi.org/10.3171/jns.1983.58.2.0270
- Nasca, R.J., McElvein, R.B., 1985. *Aspergillus fumigatus* osteomyelitis of the thoracic spine treated by excision and interbody fusion. *Spine*, **10**(9):848-850. http://dx.doi.org/10.1097/00007632-198511000-00013
- Palmisano, A., Benecchi, M., Filippo, M.D., 2011. Candida

- *sake* as the causative agent of spondylodiscitis in a hemodialysis patient. *Spine J.*, **11**(3):e12-e16. http://dx.doi.org/10.1016/j.spinee.2011.01.026
- Pappas, P.G., Kauffman, C.A., David, A., et al., 2009. Clinical practice guidelines for the management of candidiasis: 2009 update by the Infectious Diseases Society of America. Clin. Infect. Dis., 48(5):503-535. http://dx.doi.org/10.1086/596757
- Pemán, J., Jarque, I., Bosch, M., *et al.*, 2006. Spondylodiscitis caused by *Candida krusei*: case report and susceptibility patterns. *J. Clin. Microbiol.*, **44**(5):1912-1914. http://dx.doi.org/10.1128/JCM.44.5.1912-1914.2006
- Rachapalli, S.M., Malaiya, R., Mohd, T., *et al.*, 2010. Successful treatment of *Candida* discitis with 5-flucytosine and fluconazole. *Rheumatol. Int.*, **30**(11):1543-1544. http://dx.doi.org/10.1007/s00296-009-1215-x
- Shashidhar, N., Tripathy, S.K., Balasubramanian, S., *et al.*, 2014. *Aspergillus* spondylodiscitis in an immunocompetent patient following spinal anesthesia. *Orthop. Surg.*, **6**(1):72-77. http://dx.doi.org/10.1111/os.12091
- Shi, J.M., Pei, X.Y., Luo, Y., et al., 2015. Invasive fungal infection in allogeneic hematopoietic stem cell transplant recipients: single center experiences of 12 years. *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol.)*, **16**(9): 796-804.
 - http://dx.doi.org/10.1631/jzus.B1500005
- Skaf, G.S., Kanafani, Z.A., Araj, G.F., et al., 2010. Non-pyogenic infections of the spine. *Int. J. Antimicrob. Agents*, **36**(2):99-105.
 - http://dx.doi.org/10.1016/j.ijantimicag.2010.03.023
- Storm, L., Lausch, K.R., Arendrup, M.C., *et al.*, 2014. Vertebral infection with *Candida albicans* failing caspofungin and fluconazole combination therapy but successfully treated with high dose liposomal amphotericin B and flucytosine. *Med. Mycol. Case Rep.*, **6**:6-9. http://dx.doi.org/10.1016/j.mmcr.2014.07.001
- Sugar, A.M., Saunders, C., Diamond, R.D., 1990. Successful treatment of *Candida* osteomyelitis with fluconazole. A noncomparative study of two patients. *Diagn. Microbiol. Infect. Dis.*, **13**(6):517-520.
- Tan, A.C., Parker, N., Arnold, M., 2014. *Candida glabrata* vertebral osteomyelitis in an immunosuppressed patient. *Int. J. Rheuma. Dis.*, **17**(2):229-231.
- Theodoros, K., Sotirios, T., 2012. Successful treatment of azole-resistant *Candida* spondylodiscitis with high-dose caspofungin monotherapy. *Rheumatol. Int.*, **32**(9):2957-2958.
 - http://dx.doi.org/10.1007/s00296-011-2121-6
- Uyeki, T.M., Cox, N.J., 2013. Global concerns regarding novel influenza A (H7N9) virus infections. *N. Engl. J. Med.*, **368**(20):1862-1864.
- Walsh, T.J., 2008. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin. Infect. Dis.*, **46**(3):327-360. http://dx.doi.org/10.1086/525258

- Williams, R.L., Fukui, M.B., Meltzer, C.C., *et al.*, 1999. Fungal spinal osteomyelitis in the immunocompromised patient: MR findings in three cases. *Am. J. Neuroradiol.*, **20**(20):381-385.
- Yagupsky, P., 2004. Kingella kingae: from medical rarity to an emerging paediatric pathogen. *Lancet Infect. Dis.*, 4(6): 358-367.
 - http://dx.doi.org/10.1016/S1473-3099(04)01046-1
- Yang, S.G., Cao, B., Liang, L.R., *et al.*, 2012. Antiviral therapy and outcomes of patients with pneumonia caused by influenza A pandemic (H1N1) virus. *PLOS ONE*, 7(1):e29652. http://dx.doi.org/10.1371/journal.pone.0029652

List of electronic supplementary materials

Case reports or case serials related to the *Candida* or *Aspergillus* spondylosidcitis

中文概要

题 目: 念珠菌脊柱感染发生在 H7N9 治愈的患者一例报 道及文献分析念珠菌脊柱感染和曲霉菌脊柱感 染的临床区别

- **时 的**:报道在 H7N9 治愈患者中发现念珠菌脊柱感染病例,并总结念珠菌和曲霉菌脊柱感染的临床表现,处理方法和预后等不同点。
- 创新点: 首次报道在 H7N9 禽流感治愈患者中出现念珠菌 脊柱感染,经过彻底的清创融合内固定和唑类抗 真菌药治疗取得良好疗效。文献研究发现了念珠菌和曲霉菌这两大主要真菌在脊柱感染中的临床表现、处理方法和临床结局均有不同。念珠菌脊柱感染治愈率为 92.3%,而曲霉菌为 70.2%。 其可能的原因有以下四点,念珠菌血培养结果阳性率更高(41.0% vs. 24.5%),神经损害发生概率更低(25.6% vs. 43.6%),手术方式更多地选用彻底病灶清除加融合内固定(60.5% vs. 39.6%),术后多选用唑类抗真菌药。
- 方 法: 在 PubMed 和 Google Scholar 搜集念珠菌和曲霉菌脊柱感染的病例,提取患者一般信息、临床症状、感染部位、病原学结果、治疗方法、治疗结局和复发等情况进行总结分析。
- **结 论:** 念珠菌脊柱感染的治愈率相对曲霉菌脊柱感染更高,这可能与念珠菌可以早期发现,神经损害较少,更多地选用彻底病灶清除和唑类抗真菌药有关。真菌脊柱感染可以通过合适的手术和抗真菌药取得良好临床疗效。
- **关键词:** 真菌脊柱感染; H7N9 感染; 念珠菌; 曲霉菌; 综述

Fungal spondylodiscitis in a patient recovered from H7N9 virus infection: a case study and a literature review of the differences between *Candida* and *Aspergillus* spondylodiscitis

Supplemental material

Case reports or case serials related to the Candida or Aspergillus spondylosidcitis

- 1. Ackerman G, Bayley JC: Candida albicans osteomyelitis in a vertebral body previously infected with Serratia marcescens. *Spine* 1990, **15**(12):1362-1363.
- 2. Almekinders LC, Greene WB: Vertebral Candida infections. A case report and review of the literature. Clinical Orthopaedics & Related Research 1991, 267(267):174-178.
- 3. Alvarez L, Abril C: **Articular aspergillosis: case report**. *Clinical Infectious Diseases* 1995, **20**(2):457-460.
- 4. Anan S, Sanjeev R, Andrea H, Goran M, Yaw AA: **Aspergillus rib and vertebral osteomyelitis in a former intravenous drug user**. *American Journal of Medicine* 2004, **116**(116):208-209.
- 5. Anderson J, Kron IL: **Treatment of Aspergillus infection of the proximal aortic prosthetic graft with associated vertebral osteomyelitis**. *Journal of Vascular Surgery* 1984, 1(4):579-581.
- 6. Andrew H, Clifton H, Subra K: Vertebral osteomyelitis due to Candida parapsilosis in a child with Crohn disease while receiving anti-TNF therapy. *Journal of Pediatric Gastroenterology & Nutrition* 2013, **56**(4).
- 7. Barbara Alexander L, Elizabeth Dodds A, Steele MP, Perfect JR: **Aspergillus fumigatus** empyema, arthritis, and calcaneal osteomyelitis in a lung transplant patient successfully treated with posaconazole. *Journal of Clinical Microbiology* 2004, **42**(3):1376-1378.
- 8. Bridwell KH, Campbell JW, Barenkamp SJ: Surgical treatment of hematogenous vertebral Aspergillus osteomyelitis. *Spine* 1976, **15**(4):281-285.
- 9. Brown DL, Musher DM, Taffet GE: **Brown DL**, **Musher DM**, **Taffet GE**. **Hematogenously acquired Aspergillus vertebral osteomyelitis in seemingly immunocompetent drug addicts**. *Western Journal of Medicine* 1987, **147**(1):84-85.
- 10. Bruns J, Hemker T, Dahmen G: Fungal spondylitis. A case of Torulopsis glabrata and Candida tropicalis infection. *Acta Orthopaedica Scandinavica* 1986, **57**(6):563-565.
- 11. Byrd BF, Weiner MH, Mcgee ZA: Aspergillus spinal epidural abscess. *Jama* 1982, **248**(248):3138-3139.
- 12. Castelli C, Benazzo F, Minoli L, Marone P, Seghezzi R, Carlizzi CN: Aspergillus infection of the L3-L4 disc space in an immunosuppressed heart transplant patient. *Spine* 1976, 15(12):1369-1373.
- 13. Chang HM, Yu HH, Yang YH, Lee WI, Lee JH, Wang LC, Lin YT, Chiang BL: Successful treatment of Aspergillus flavus spondylodiscitis with epidural abscess in a patient with chronic granulomatous disease. *Pediatric Infectious Disease Journal* 2012, 31(1):100-101.
- 14. Chi CY, Fung CP, Liu CY: Aspergillus flavus epidural abscess and osteomyelitis in a diabetic patient. *Journal of microbiology, immunology, and infection* 2003, **36**(2):145-148.
- 15. Chia SL, Tan BH, Tan CT, Tan SB: Candida spondylodiscitis and epidural abscess:

- management with shorter courses of anti-fungal therapy in combination with surgical debridement. *Journal of Infection* 2005, **51**(1):17-23.
- 16. Christelle D-L, Claire L, Frédérque C-G, Lucia P, Emmanuelle D: **Candida albicans diskitis after body piercing in an immunocompetent patient**. *Joint Bone Spine* 2013, **80**(2):226–227.
- 17. Cone LA, Byrd RG, Potts BE, Moxell W: Diagnosis and treatment of Candida vertebral osteomyelitis: clinical experience with a short course therapy of amphotericin B lipid complex. Surgical neurology 2004, 62(3):234–237.
- 18. Cortet B, Richard R, Deprez X, Lucet L, Flipo RM, Le Loët X, Duquesnoy B, Delcambre B: Aspergillus spondylodiscitis: successful conservative treatment in 9 cases. *Journal of Rheumatology* 1994, **21**(7):1287-1291.
- 19. Derkinderen P, Bruneel F, Bouchaud O, Regnier B: **Spondylodiscitis and epidural abscess due to Candida albicans**. *European spine journal* 2000, **9**(1):72-74.
- 20. D'Hoore K, Hoogmartens M: Vertebral aspergillosis. A case report and review of the literature. Acta Orthopaedica Belgica 1993, 59(3):306-314.
- 21. Dietz R, Huber G, Thetter O, Volkmer I: **Aspergillosis of the lung with osteoclasis and paraplegia**. *Neuroradiology* 1982, **23**(4):219-221.
- Dijkmans BAC, Broek PJVD, Meer JWMVD, Koolen MI, Mouton RP, Falke THM: Hematogenous candida vertebral osteomyelitis treated with ketoconazole. *Infection* 1982, 10(5):290-292.
- 23. Edwards JE, Turkel SB, Elder HA, Rand RW, Guze LB: **Hematogenous candida** osteomyelitis. Report of three cases and review of the literature. *American Journal of Medicine* 1975, **59**(1):89–94.
- 24. Eisen DP, Macginley R, Christensson B, Larsson L, Woods ML: Candida tropicalis vertebral osteomyelitis complicating epidural catheterisation with disease paralleled by elevated D-arabinitol/L-arabinitol ratios. European Journal of Clinical Microbiology 2000, 19(1):61-63.
- 25. Eismont FJ, Bohlman HH, Soni PL, Goldberg VM, Freehafer AA: Pyogenic and fungal vertebral osteomyelitis with paralysis. *Journal of Bone & Joint Surgery* 1983, **65**(1):19-29.
- 26. Ersoy A, Akdag I, Akalin H, Sarisozen B, Ener B: **Aspergillosis osteomyelitis and joint infection in a renal transplant recipient**. *Transplantation Proceedings* 2007, **39**(5):1662–1663.
- 27. Ersoy A, Dizdar OS, Koc AO: Aspergillus fumigatus spondylodiskitis in renal transplant patient: voriconazole experience. Experimental & Clinical Transplantation 2011, 9(4):265-269.
- 28. Ferra C, Gingrich RD: Candida tropicalis vertebral osteomyelitis: A late sequela of fungemia. Clinical Infectious Diseases 1994, 19(4):697-703.
- 29. Frazier DD, Campbell DR, Garvey TA, Wiesel S, Bohlman HH, Eismont FJ: Fungal infections of the spine. Report of eleven patients with long-term follow-up. *Journal of Bone & Joint Surgery American Volume* 2001, 83-a(83-A):560-565.
- 30. Friedman BC, Simon GL: Candida vertebral osteomyelitis: report of three cases and a review of the literature. *Diagnostic Microbiology & Infectious Disease* 1987, 8(1):31–36.
- 31. Gathe JC, Jr., Harris RL, Garland B, Jr.: Candida osteomyelitis. Report of five cases and review of the literature. *American Journal of Medicine* 1987, **82**(5):927–937.

- 32. Glotzbach RE: Aspergillus terreus infection of pseudoaneurysm of aortofemoral vascular graft with contiguous vertebral osteomyelitis. *American journal of clinical pathology* 1982, 77(2):224-227.
- 33. Govender S, Rajoo R, Goga IE, Charles RW: **Aspergillus osteomyelitis of the spine**. *Spine* 1991, **16**(7):746-749.
- 34. Grandiere-Perez L, Asfar P, Foussard C, Chennebault J, Penn P, Degasne I: **Spondylodiscitis** due to Aspergillus terreus during an efficient treatment against invasive pulmonary aspergillosis. *Intensive Care Medicine* 2000, 26(7):1010-1011.
- 35. Grossman M: Aspergillosis of bone. British Journal of Radiology 1975, 48(565):57-59.
- 36. Gupta PK, Mahapatra AK, Gaind R, Bhandari S, Musa MM, Lad SD: **Aspergillus spinal epidural abscess**. *Pediatric Neurosurgery* 2001, **35**(1):18-23.
- 37. Mouas H, Lutsar I, Dupont B, Fain O, Herbrecht R, Lescure F, Lortholary O: Voriconazole for invasive bone aspergillosis: a worldwide experience of 20 cases. *Clinical Infectious Diseases* 2005, 40(8):1141-1147.
- 38. Hendrickx L, Wijngaerden EV, Samson I, Peetermans WE: Candidal vertebral osteomyelitis: report of 6 patients, and a review. Clinical Infectious Diseases 2001, 32(4):527-533.
- 39. Hennequin C, Bourée P, Hiesse C, Dupont B, Charpentier B: **Spondylodiskitis due to** Candida albicans: report of two patients who were successfully treated with fluconazole and review of the literature. *Clinical Infectious Diseases* 1996, **23**(1):176-178.
- 40. Herzog W, Perfect J, Roberts L: **Intervertebral diskitis due to Candida tropicalis**. *Southern medical journal* 1989, **82**(2):270-273.
- 41. Hirschmann JV, Everett ED: **Candida vertebral osteomyelitis**. *Journal of Bone & Joint Surgery American Volume* 1976, **58**(4):573-575.
- 42. Holmes PF, Osterman DW, Tullos HS: **Aspergillus discitis. Report of two cases and review of the literature**. *Clinical Orthopaedics & Related Research* 1988, **226**(226):240-246.
- 43. Hummel M, Schüler S, Weber U, Schwertlick G, Hempel S, Theiss D, Rees W, Mueller J, Hetzer R: Aspergillosis with Aspergillus osteomyelitis and diskitis after heart transplantation: surgical and medical management. *Journal of Heart & Lung Transplantation* 1993, 12(4):599-603.
- 44. Hung CC, Chang SC, Yang PC, Hsieh WC: Invasive pulmonary pseudallescheriasis with direct invasion of the thoracic spine in an immunocompetent patient. European Journal of Clinical Microbiology & Infectious Diseases 1994, 13(9):749-751.
- 45. Ingwer I, Mcleish KR, Tight RR, White AC: **Aspergillus fumigatus epidural abscess in a renal transplant recipient**. *Archives of Internal Medicine* 1978, **138**(1):153-153.
- 46. Iwata A, Ito M, Abumi K, Sudo H, Kotani Y, Shono Y, Minami A: Fungal Spinal Infection Treated with Percutaneous Posterolateral Endoscopic Surgery. Central European Neurosurgery 2014, 75(3):170-176.
- 47. Seres JL, Ono H, Benner EJ: **Aspergillosis presenting as spinal cord compression. Case report**. *Journal of Neurosurgery* 1972, **36**(2):221-224.
- 48. Javier P, Isidro J, María B, Emilia C, Miguel S, Rosario DL, Araceli M: **Spondylodiscitis** caused by Candida krusei: case report and susceptibility patterns. *Journal of Clinical Microbiology* 2006, 44(5):1912-1914.
- 49. Jiang Z, Wang Y, Jiang Y, Xu Y, Meng B: Vertebral osteomyelitis and epidural abscess due to Aspergillus nidulans resulting in spinal cord compression: Case report and literature

- review. Journal of International Medical Research 2013, 41(2):502-510.
- Jonnalagadda S, Veerabagu MP, Rakela J, Kusne S, Randhawa P, Rabinovitz M: Candida albicans osteomyelitis in a liver transplant recipient: a case report and review of the literature. *Transplantation* 1996, **62**(8):1182-1184.
- Jorge G, Isabelle S, Daniel L, Kalthum BW, Peter R: An unusual cause of vertebral osteomyelitis: Candida species. Scandinavian Journal of Infectious Diseases 2003, 35(4):288-291.
- 52. Jorge VC, Cardoso C, Noronha C, Simões J, Riso N, Vaz RM: Fungal spondylodiscitis in a non-immunocompromised patient. *Bmj Case Reports* 2012, **2012**:bcr1220115337.
- 53. Karthik K, Shetty AP, Rajasekaran S: **Spontaneous cord transection due to invasive aspergillus spondylitis in an immunocompetent child**. *European Spine Journal* 2011, **20**(2 Supplement):188-192.
- 54. Kashimoto T, Kitagawa H, Kachi H: Candida Tropicalis Vertebral Osteomyelitis and Discitis. Spine 1986, 11(1):57-61.
- 55. Kelesidis T, Tsiodras S: Successful treatment of azole-resistant Candida spondylodiscitis with high-dose caspofungin monotherapy. Rheumatology International 2012, 32(9):2957-2958.
- 56. Kolbe ABL, Mckinney AM, Kendi ATK, Misselt D: Aspergillus meningitis and discitis from low-back procedures in an immunocompetent patient. *Acta radiologica* 2007, 48(6):687-689.
- 57. Lang EW, Pitts LH: Intervertebral disc space infection caused by Aspergillus fumigatus. *European Spine Journal* 1996, **5**(3):207-209.
- 58. Langlois RP, Flegel KM, Meakins JL, Morehouse DD, Robson HG, Guttmann RD: Cutaneous aspergillosis with fatal dissemination in a renal transplant recipient. Canadian Medical Association journal 1980, 122(6):673-676.
- 59. Laurent S, Daniel VL, Christian B, Antoine DT, Oscar M, Francois P, Daniel G: Candida glabrata spinal osteomyelitis involving two contiguous lumbar vertebrae: a case report and review of the literature. Diagnostic Microbiology & Infectious Disease 2003, 45(2):137-141.
- 60. Lenzi J, Agrillo A, Santoro A, Marotta N, Cantore GP: **Postoperative spondylodiscitis from Aspergillus fumigatus in immunocompetent subjects**. *Journal of Neurosurgical Sciences* 2004, **48**(2):81-85.
- 61. Parry MF, Grant B, Yukna M, Adler-Klein D, McLeod GX, Taddonio R, Rosenstein C: Candida osteomyelitis and diskitis after spinal surgery: an outbreak that implicates artificial nail use. Clinical Infectious Diseases 2001, 32(3):352-357.
- 62. Martinez M, Lee AS, Hellinger WC, Kaplan J: Vertebral Aspergillus Osteomyelitis and Acute Diskitis in Patients With Chronic Obstructive Pulmonary Disease. *Mayo Clinic Proceedings Mayo Clinic* 1999, 74(6):579-583.
- 63. Mary Jane B, Pratik S, Edwin S: Misidentification of Candida parapsilosis as C famata in a clinical case of vertebral osteomyelitis. American Journal of the Medical Sciences 2011, 341(1):71-73.
- 64. Mawk JR, Erickson DL, Chou SN, Seljeskog EL: **Aspergillus infections of the lumbar disc spaces. Report of three cases**. *Journal of Neurosurgery* 1983, **58**(2):270-274.
- 65. Mcgregor A, Mcnicol D, Collignon P: Aspergillus-induced discitis. A role for itraconazole

- in therapy? Spine 1976, 17(12):1512-1514.
- 66. Mckee DF, Barr WM, Bryan CS, Lunceford EM: **Primary aspergillosis of the spine** mimicking Pott's paraplegia. *Journal of Bone & Joint Surgery American Volume* 1985, **66**(9):1481-1483.
- 67. Miller DJ, Mejicano GC: Vertebral osteomyelitis due to Candida species: case report and literature review. Clinical Infectious Diseases 2001, 33(4):523-530.
- 68. Morgenlander JC, Rossitch E, Rawlings CE: **Aspergillus disc space infection: case report and review of the literature**. *Neurosurgery* 1989, **25**(1):126-129.
- 69. Mullins RF, Jr JMS, Savage J, Davis JB, Law EJ: Osteomyelitis of the spine in a burn patient due to Candida albicans. Burns Journal of the International Society for Burn Injuries 1993, 19(2):174–176.
- 70. Munk PL, Lee MJ, Poon PY, O'Connell JX, Coupland DB, Janzen DL, Logan PM, Dvorak MF: Candida osteomyelitis and disc space infection of the lumbar spine. *Skeletal Radiology* 1997, **26**(1):42-46.
- 71. Nandeesh BN, Usha K, Betty A: Vertebral osteomyelitis with a rare etiology diagnosed by fine-needle aspiration cytology. *Diagnostic Cytopathology* 2010, **38**(5):360-363.
- 72. Nandeesh S, Sujit Kumar T, Sakthivel B, Naresh D, Sathyanarayana V: Aspergillus spondylodiscitis in an immunocompetent patient following spinal anesthesia. *Orthopaedic surgery* 2014, 6(1):72-77.
- 73. Nasca RJ, Mcelvein RB: Aspergillus fumigatus osteomyelitis of the thoracic spine treated by excision and interbody fusion. *Spine* 1976, **10**(9):848-850.
- 74. Sinthumathi N, Georgi A, Milly M, Lalitha MK, Srinivasan CN: Secondary sternal Aspergillus osteomyelitis in a diabetic hemodialysis patient with previous allograft rejection. *Hemodialysis International* 2007, 11(4):403–405.
- 75. Neale TJ, Muir JC, Mills H, Horne JG, Jones MR: Candida albicans vertebral osteomyelitis in chronic renal failure. *Postgraduate Medical Journal* 1987, **63**(742):1059-1060.
- 76. Nicolle A, Blanchardière A, De La, Bonhomme J, Hamon M, Leclercq R, Hitier M: Aspergillus vertebral osteomyelitis in immunocompetent subjects: case report and review of the literature. *Infection* 2013, 41(4):833-840.
- 77. Oh IS, Seo JY, Ha KY, Kim YC: **Treatment for Multiple Aspergillus Spondylitis Including a Hip Joint**. *Asian Spine Journal* 2009, **3**(2):106–112.
- 78. Oksi J, Finnilä T, Hohenthal U, Rantakokko-Jalava K: Candida dubliniensis spondylodiscitis in an immunocompetent patient. Case report and review of the literature. *Medical Mycology Case Reports* 2014, 3:4-7.
- 79. Ooij AV, Beckers JM, Herpers MJ, Walenkamp GH: Surgical treatment of aspergillus spondylodiscitis. European Spine Journal 2000, 9(1):75-79.
- 80. Korovessis P, Repanti M, Katsardis T, Stamatakis M: Anterior decompression and fusion for Aspergillus osteomyelitis of the lumbar spine associated with paraparesis. *Spine* 1994, 19(23):2715-2718.
- 81. Palmisano A, Benecchi M, Filippo MD: Candida sake as the causative agent of spondylodiscitis in a hemodialysis patient. *The Spine Journal* 2011, 11(3):e12–e16.
- Park KU, Lee HS, Kim CJ, Kim EC: Fungal discitis due to Aspergillus terreus in a patient with acute lymphoblastic leukemia. *Journal of Korean Medical Science* 2000, 15(6):704-707.

- 83. Park SB, Kang MJ, Whang EA, Han SY, Kim HC: A case of fungal sepsis due to aspergillus spondylitis followed by cytomegalovirus infection in a renal transplant recipient. *Transplantation Proceedings* 2004, **36**(7):2154-2155.
- 84. Pennisi AK, Davis DO, Wiesel S, Moskovitz P: **CT appearance of Candida diskitis**. *Journal of Computer Assisted Tomography* 1985, **9**(6):1050-1054.
- 85. Peters-Christodoulou MN, de Beer FC, Bots GT, Ottenhoff TM, Thompson J, Van 'T Wout J: Treatment of postoperative Aspergillus fumigatus spondylodiscitis with itraconazole. Scandinavian Journal of Infectious Diseases 1991, 23(3):373-376.
- 86. Pohjola-Sintonen S, Ruutu P, Tallroth K: **Hematogenous Candida spondylitis. A case report**. *Acta Medica Scandinavica* 1984, **215**(1):85–87.
- 87. Rachapalli SM, Malaiya R, Mohd T, Hughes RA: Successful treatment of Candida discitis with 5-flucytosine and fluconazole. *Rheumatology International* 2010, 30(11):1543-1544.
- 88. Ranjan R, Mishra S, Ranjan S: **Aspergillus vertebral osteomyelitis in an immunocompetent person**. *Neurology India* 2010, **58**(5):806-807.
- 89. Redmond A, Carré IJ, Biggart JD, Mackenzie DWR: **Aspergillosis (Aspergillus nidulans)** involving bone. *Journal of Pathology & Bacteriology* 1965, **89**(1):391-395.
- 90. Rhame FS, Brown B, Thompson RJ: **Aspergillus osteomyelitis: Report of four cases and review of the literature**. *American Journal of Medicine* 1982, **73**(2):295–300.
- 91. Rössel P, Schønheyder HC, Nielsen H: **Fluconazole therapy in Candida albicans spondylodiscitis**. *Infectious Diseases* 1998, **30**(5):527-530.
- 92. Brandt SJ, Thompson RL, Wenzel RP: Mycotic pseudoaneurysm of an aortic bypass graft and contiguous vertebral osteomyelitis due to Aspergillus fumigatus. *American Journal of Medicine* 1985, **79**(79):259-262.
- 93. Salvalaggio PRO, Bassetti M, Lorber MI, Micheletto GC, Friedman AL, Andriole VT, Basadonna GP: **Aspergillus vertebral osteomyelitis after simultaneous kidney-pancreas transplantation**. *Transplant Infectious Disease* 2003, **5**(5):187-190.
- 94. Savall F, Dedouit F, Telmon N, Rougé D: Candida albicans spondylodiscitis following an abdominal stab wound: forensic considerations. *Journal of Forensic & Legal Medicine* 2014, **23**(2):1-3.
- 95. Schilling A, Seibold M, V, Gleissner B: Successfully treated Candida krusei infection of the lumbar spine with combined caspofungin/posaconazole therapy. *Medical Mycology* 2008, **46**(1):79-83.
- 96. Schubert M, Schär G, Curt A, Dietz V: Aspergillus spondylodiscitis in an immunocompetent paraplegic patient. Spinal cord 1998, 36(11):800-803.
- 97. Sebastiani GD, Galas F: **Spondylodiscitis due to Candida tropicalis as a cause of inflammatory back pain**. Clinical Rheumatology 2001, **20**(6):435-437.
- 98. Shaikh BS, PhD PCAM, Aber RC: Vertebral disc space infection and osteomyelitis due to candida albicans in a patient with acute myelomonocytic leukemia. *Cancer* 1980, 45(5):1025–1028.
- 99. Shaw FW, Warthen HJ: **Aspergillosis of Bone**. Southern medical journal 1936, **29**(11):1070-1071.
- 100. Son JM, Jee WH, Jung CK, Kim SI, Ha KY: Aspergillus spondylitis involving the cervico-thoraco-lumbar spine in an immunocompromised patient: a case report. *Korean Journal of Radiology* 2007, **8**(5):448-451.

- 101. Storm L, Lausch KR, Arendrup MC, Mortensen KL, Petersen E: Vertebral infection with Candida albicans failing caspofungin and fluconazole combination therapy but successfully treated with high dose liposomal amphotericin B and flucytosine. *Medical Mycology Case Reports* 2014, 6:6-9.
- 102. Stratov I, Korman TM, Johnson PDR: Management of Aspergillus osteomyelitis: report of failure of liposomal amphotericin B and response to voriconazole in an immunocompetent host and literature review. European Journal of Clinical Microbiology & Infectious Diseases 2003, 22(5):277-283.
- 103. Studemeister A, Stevens DA: Aspergillus Vertebral Osteomyelitis in Immunocompetent Hosts: Role of Triazole Antifungal Therapy. Clinical Infectious Diseases 2011, 52(1):e1-6.
- 104. Sugar AM, Saunders C, Diamond RD: Successful treatment of Candida osteomyelitis with fluconazole. A noncomparative study of two patients. *Diagnostic Microbiology & Infectious Disease* 1990, **13**(6):517–520.
- Taillandier J, Alemanni M, Cerrina J, Ladurie FLR, Dartevelle P: **Aspergillus osteomyelitis** after heart-lung transplantation. *Journal of Heart & Lung Transplantation* 1997, 16(4):436-438.
- Takagi K, Yoshida A, Yamauchi T, Yamashita T, Iwasaki H, Tsutani H, Maezawa Y, Baba H, Ueda T: Successful treatment of Aspergillus spondylodiscitis with high-dose itraconazole in a patient with acute myelogenous leukemia. *Leukemia* 2001, **15**(10):1670-1671.
- 107. Tan AC, Parker N, Arnold M: Candida glabrata vertebral osteomyelitis in an immunosuppressed patient. International Journal of Rheumatic Diseases 2014, 17(2):229–231.
- 108. Tang TJ, Janssen HL, Ch VDV, de Man RA, Metselaar HJ, Tilanus HW, De MS: Aspergillus osteomyelitis after liver transplantation: conservative or surgical treatment? European Journal of Gastroenterology & Hepatology 2000, 12(1):123-126.
- 109. Tew CW, Han FC, Jureen R, Tey BH: **Aspergillus vertebral osteomyelitis and epidural abscess**. *Singapore medical journal* 2009, **50**(4):437, e151-e154.
- 110. Turner DL, Johnson SA, Rule SA: Successful treatment of candidal osteomyelitis with fluconazole following failure with liposomal amphotericin B. *Journal of Infection* 1999, 38(1):51-53.
- 111. Vaishya S, Sharma MS: Spinal aspergillus vertebral osteomyelitis with extradural abscess: case report and review of literature. Surgical neurology 2004, 61(6):551–555.
- 112. Vinas F, King P, Diaz F: **Spinal aspergillus osteomyelitis**. *Clinical Infectious Diseases* 1999, **28**(6):1223-1229.
- 113. Assaad W, Nuchikat PS, Cohen L, Esguerra JV, Whittier FC: Aspergillus discitis with acute disc abscess. *Spine* 1976, **19**(19):2226-2229.
- 114. Wéclawiak H, Garrouste C, Kamar N, Linas M-D, Tall P, Dambrin C, Durand D, Rostaing L: Aspergillus fumigatus-related spondylodiscitis in a heart transplant patient successfully treated with voriconazole. *Transplantation Proceedings* 2007, 39(8):2627–2628.
- 115. Werner BC, Hogan MV, Shen FH: Candida lusitaniae discitis after discogram in an immunocompetent patient. Spine Journal 2011, 11(10):e1-e6.
- Williams RL, Fukui MB, Meltzer CC, Swarnkar A, Johnson DW, Welch W: Fungal spinal osteomyelitis in the immunocompromised patient: MR findings in three cases. *American Journal of Neuroradiology* 1999, **20**(20):381-385.

- 117. Witzig RS, Greer DL, Jr HN: **Aspergillus flavus mycetoma and epidural abscess successfully treated with itraconazole**. *Journal of medical and veterinary mycology* 1996, **34**(2):133-137.
- 2 Zhu LP, Chen XS, Wu JQ, Yang FF, Weng XH: Aspergillus vertebral osteomyelitis and ureteral obstruction after liver transplantation. Transplant Infectious Disease 2011, 13(2):192-199.